

Why Patients Visit Their Doctors: Assessing the Most Prevalent Conditions in a Defined American Population

Jennifer L. St. Sauver, PhD, MPH; David O. Warner, MD; Barbara P. Yawn, MD, MSc; Debra J. Jacobson, MS; Michaela E. McGree, BS; Joshua J. Pankratz, BS; L. Joseph Melton III, MD, MPH; Véronique L. Roger, MD, MPH; Jon O. Ebbert, MD; and Walter A. Rocca, MD, MPH

Abstract

Objective: To describe the prevalence of nonacute conditions among patients seeking health care in a defined US population, emphasizing age, sex, and ethnic differences.

Patients and Methods: The Rochester Epidemiology Project (REP) medical records linkage system was used to identify all residents of Olmsted County, Minnesota, on April 1, 2009, who had consented to review of their medical records for research (142,377 patients). We then electronically extracted all *International Classification of Diseases, Ninth Revision* codes noted in the records of these patients by any health care institution between January 1, 2005, and December 31, 2009. We grouped *International Classification of Diseases, Ninth Revision* codes into clinical classification codes and then into 47 broader disease groups associated with health-related quality of life. Age- and sex-specific prevalence was estimated by dividing the number of individuals within each group by the corresponding age- and sex-specific population. Patients within a group who had multiple codes were counted only once.

Results: We included a total of 142,377 patients, 75,512 (53%) of whom were female. Skin disorders (42.7%), osteoarthritis and joint disorders (33.6%), back problems (23.9%), disorders of lipid metabolism (22.4%), and upper respiratory tract disease (22.1%, excluding asthma) were the most prevalent disease groups in this population. Ten of the 15 most prevalent disease groups were more common in women in almost all age groups, whereas disorders of lipid metabolism, hypertension, and diabetes were more common in men. Additionally, the prevalence of 7 of the 10 most common groups increased with advancing age. Prevalence also varied across ethnic groups (whites, blacks, and Asians).

Conclusion: Our findings suggest areas for focused research that may lead to better health care delivery and improved population health.

© 2013 Mayo Foundation for Medical Education and Research ■ Mayo Clin Proc. 2013;88(1):56-67



From the Division of Epidemiology (J.L.S., B.P.Y., L.J.M., V.L.R., W.A.R.), Department of Anesthesiology (D.O.W.), Division of Biomedical Statistics and Informatics (D.J.J., M.E.M.), Division of Biomedical Informatics Support System (J.J.P.), Division of Cardiovascular Diseases (V.L.R.), Division of Primary Care Internal Medicine (J.O.E.), and Department of Neurology (W.A.R.), Mayo Clinic, Rochester, MN; and Department of Research, Olmsted Medical Center, Rochester, MN (B.P.Y.).

Chronic diseases account for the majority of health care utilization and expenditures in middle-aged and older populations.¹⁻³ As the population ages, more individuals are living with multiple chronic medical conditions. One-fourth of Americans with chronic conditions account for almost two-thirds of the total US health care expenditures.⁴ Research on chronic disease has largely focused on a specific group of conditions with high morbidity and mortality (including diabetes and chronic heart disease). However, other types of nonacute conditions, with less severe long-term outcomes, may affect large segments of the population and may

account for a substantial amount of health care resource utilization. Recognition of these other conditions may suggest new areas for improving health care delivery and population health management.

Health care reform has intensified the need for information on health care resource utilization for nonacute conditions. The Patient Protection and Affordable Care Act allows the restructuring of Medicare reimbursements into “bundled payments.”⁵ This restructuring will require the rational deployment of treatment resources to ensure the financial solvency of medical institutions. Additionally, clinical decision support for chronic diseases has

been identified as critical for the patient-centered medical home model.⁶ However, development of these models requires quantification of prevalent chronic diseases across populations.

Unfortunately, the prevalence of diseases can be difficult to capture across all age groups because only a few databases in the United States include younger populations. Additionally, it can be difficult to consider the prevalence of multiple conditions concurrently in a single population. Failure to simultaneously consider all possible drivers of health care utilization can result in inefficient targeting of resources to improve population health.

To address these problems, we conducted a study to identify the prevalence of the most common nonacute conditions in a defined US population using the resources of the Rochester Epidemiology Project (REP). The REP medical records linkage system provides an ideal opportunity to quantify the prevalence of all medical conditions in an entire population, across age, sex, and ethnic groups, regardless of socioeconomic or insurance status.⁷

PATIENTS AND METHODS

Study Population

The REP links data on medical care delivered to the population of Olmsted County, Minnesota.⁷⁻⁹ The vast majority of medical care in this community is currently provided by a few health care institutions: the Mayo Clinic and its 2 affiliated hospitals, Olmsted Medical Center and its affiliated hospital, and the Rochester Family Medicine Clinic. The health care records from these institutions are linked together through the REP records linkage system.^{8,9} Patients are categorized as residents or nonresidents of Olmsted County at the time of each health care visit on the basis of their address. Over the years, this address information has been accumulated and is used in medical research to determine the number of residents in Olmsted County at any given point in time since 1966 (the REP Census). The population counts obtained by the REP Census are similar to those obtained by the US Census, indicating that virtually the entire population of the county is captured by the system.^{8,9} For this study, we used the REP Census to identify all individuals who resided in Olmsted County on April 1,

2009, but we excluded those individuals who had not given permission to at least one health care institution for use of their medical records for research.⁸

Definition of Disease Groups

The diagnostic indices of the REP were searched electronically to extract all *International Classification of Diseases, Ninth Revision (ICD-9)* codes in the medical records of members of the Olmsted County population assigned by any health care institution from January 1, 2005, through December 31, 2009. These ICD-9 codes were first grouped into clinical classification codes (CCCs) proposed by the Agency for Healthcare Research and Quality-Healthcare Cost and Utilization Project.^{10,11} For this study, we focused specifically on conditions that were not likely to resolve in a short period of time and that were likely to require multiple health care visits over several years for evaluation and treatment. However, these conditions were not confined to conditions typically considered chronic diseases, such as diabetes and heart disease. For example, we included conditions such as tuberculosis, back problems, and esophageal disorders. We excluded conditions related to dental or vision problems because the REP does not capture all data from local dentists or optometrists. These CCCs were then combined into broader disease groups that have been associated with health-related quality of life, such as cancer, diabetes, thyroid disorders, and heart failure, according to the classification system developed by Mukherjee et al.^{11,12} We modified this system by using updated CCCs and by including breast, uterine, ovarian, and prostate cancer in the cancer category, but excluding benign neoplasms and neoplasms of uncertain malignancy.¹² The final CCCs used for this study and the modified disease groups are shown in Supplemental Table 1 (available online at <http://www.mayoclinicproceedings.org>).

Statistical Analyses

The point prevalence of each CCC was measured using April 1, 2009, as the prevalence day.¹³ The history of a given disease on the prevalence day was derived from a 5-year capture time frame (the 5 years preceding the prevalence day). In general, for nonacute

conditions, our findings should be comparable to point prevalence figures derived from a population survey.¹³ The crude age- and sex-specific prevalence of each of the 47 disease groups was estimated by dividing the number of individuals in a group by the corresponding age- and sex-specific Olmsted County population on the prevalence day. These prevalence figures were directly standardized to the 2000 total US population by age and by sex when appropriate to make comparisons of aggregated data (2000 US Census). This study covered the target population completely, and no sampling was involved. For this reason, statistical tests may not be appropriate, and confidence intervals were not included in the tables.¹⁴⁻¹⁶

RESULTS

Description of the Olmsted County Population

Overall, the REP infrastructure captured 146,687 Olmsted County residents in 2009 compared with 143,962 individuals predicted by the US Census.¹⁷ Therefore, the REP captured slightly more people than the US Census (101.9%). These results are consistent with a previous study that examined REP capture rates between 1970 and 2000.⁸

Of 146,687 residents, 142,377 gave permission for use of their medical records for research (97.1%). The population included 75,512 females (53%). Age and sex distributions were virtually identical to US Census estimates.¹⁸ However, the proportion of people in the white category was lower, and the proportion in the other/unknown ethnic category was higher compared with US Census estimates. Because we presume that most of the patients in the other/unknown category were white (85.7% of the population self-reported white race in the 2010 census), we grouped the other/unknown category with the white category.

Results by Broad Disease Groups

Table 1 shows the 20 most prevalent conditions. Data for the remaining 27 disease groups are shown in Supplemental Table 2 (available online at <http://www.mayoclinicproceedings.org>). Skin disorders were the most prevalent disease group in this population. Almost half of the population (42.7%)

had at least one ICD-9 code for a skin condition within approximately 5 years. Skin disorders were followed in frequency by osteoarthritis and joint disorders (33.6%), back problems (23.9%), disorders of lipid metabolism (22.4%), and upper respiratory tract disease (22.1%). By contrast, systemic lupus erythematosus and connective tissue disorders, tuberculosis, human immunodeficiency virus infection, sickle cell anemia, and cystic fibrosis were the least prevalent conditions (Supplemental Table 2). Seven of the 10 most prevalent disease groups increased with advancing age (Table 1). However, the prevalence of upper respiratory tract disease remained relatively consistent across all age groups. The prevalence of anxiety, depression, and bipolar disorders was low in 0- to 18-year-olds, increased dramatically in 19- to 29-year-olds, and remained constant across the older age groups. Headaches, including migraine, also increased in the 19- to 29-year-olds but declined after age 50 years.

The most prevalent disease groups differed by age. For example, skin disorders were the most prevalent condition in 0 to 18-year-olds, followed by upper respiratory tract disease and osteoarthritis and joint disorders. By contrast, hypertension was the most prevalent condition in patients who were 65 years or older, followed by disorders of lipid metabolism and skin disorders (Figure 1). Ten of the 15 most prevalent disease groups were more common in women in almost all age groups, whereas disorders of lipid metabolism, hypertension, and diabetes were more common in men (Figure 2).

The prevalence of the top 10 disease groups also differed by ethnic group (Figure 3). Blacks had a higher prevalence for 7 of the top 10 disease groups. The biggest differences in blacks vs whites were a higher prevalence of back problems and headaches, including migraine, in blacks. In contrast, whites had a higher prevalence of skin disorders compared with both blacks and Asians. Asians had a higher prevalence of diabetes than whites.

Results for Specific ICD-9 Codes

Although our primary analyses considered a high-level grouping of diseases, we also examined the individual ICD-9 codes within the groups. The prevalence estimates of selected

TABLE 1. Age- and Sex-Specific Prevalence (per 100 Population) of the 20 Most Common Chronic Disease Groups in the 2009 Olmsted County, Minnesota. Population (N=142,377)^a

Chronic disease group	Age (y)										All ages			
	0-18		19-29		30-49		50-64		≥65		Crude ^b		Standardized ^c	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Skin disorders														
Both sexes	12,703	32.95	9170	38.26	15,652	41.27	12,390	50.39	11,398	65.75	61,313	43.06	61,313	42.67
Men	6232	31.78	3247	31.41	5923	33.11	5221	45.42	4980	66.11	25,603	38.29	25,603	38.43
Women	6471	34.15	5923	43.45	9729	48.55	7169	54.76	6418	65.47	35,710	47.29	35,710	46.90
Osteoarthritis and joint disorders														
Both sexes	5580	14.47	6044	25.22	13,122	34.60	12,275	49.92	10,971	63.28	47,992	33.71	47,992	33.58
Men	2859	14.58	2752	26.62	5832	32.60	5223	45.43	4273	56.72	20,939	31.32	20,939	31.72
Women	2721	14.36	3292	24.15	7290	36.38	7052	53.87	6698	68.33	27,053	35.83	27,053	35.13
Back problems														
Both sexes	2193	5.69	4890	20.40	11,054	29.15	8287	33.70	7692	44.37	34,116	23.96	34,116	23.90
Men	1050	5.35	1653	15.99	4588	25.65	3508	30.52	2966	39.37	13,765	20.59	13,765	21.12
Women	1143	6.03	3237	23.75	6466	32.27	4779	36.50	4726	48.21	20,351	26.95	20,351	26.48
Disorders of lipid metabolism														
Both sexes	135	0.35	704	2.94	7261	19.15	11,948	48.59	12,143	70.05	32,191	22.61	32,191	22.39
Men	75	0.38	330	3.19	4247	23.74	6110	53.15	5463	72.52	16,225	24.27	16,225	24.74
Women	60	0.32	374	2.74	3014	15.04	5838	44.59	6680	68.14	15,966	21.14	15,966	20.19
Other upper respiratory tract disease														
Both sexes	9184	23.82	4597	19.18	8436	22.24	5339	21.71	3941	22.73	31,497	22.12	31,497	22.10
Men	5033	25.66	1765	17.08	3470	19.40	2221	19.32	1717	22.79	14,206	21.25	14,206	21.16
Women	4151	21.91	2832	20.78	4966	24.78	3118	23.82	2224	22.69	17,291	22.90	17,291	22.99
Anxiety, depression, and bipolar disorders														
Both sexes	2559	6.64	5577	23.27	9927	26.17	6127	24.92	4156	23.97	28,346	19.91	28,346	19.75
Men	1179	6.01	1775	17.17	3453	19.30	2139	18.61	1346	17.87	9892	14.79	9892	15.09
Women	1380	7.28	3802	27.89	6474	32.31	3988	30.46	2810	28.67	18,454	24.44	18,454	24.13
Chronic neurologic disorders														
Both sexes	2774	7.19	2812	11.73	7482	19.73	6829	27.77	8324	48.02	28,221	19.82	28,221	19.75
Men	1519	7.75	995	9.63	2929	16.37	2894	25.17	3412	45.29	11,749	17.57	11,749	17.92
Women	1255	6.62	1817	13.33	4553	22.72	3935	30.06	4912	50.11	16,472	21.81	16,472	21.43
Hypertension														
Both sexes	108	0.28	513	2.14	4450	11.73	8918	36.27	12,251	70.67	26,240	18.43	26,240	18.21
Men	64	0.33	269	2.60	2444	13.66	4514	39.27	5290	70.22	12,581	18.82	12,581	19.22
Women	44	0.23	244	1.79	2006	10.01	4404	33.64	6961	71.01	13,659	18.09	13,659	17.22
Headaches, including migraines														
Both sexes	3286	8.52	4135	17.25	6753	17.81	3745	15.23	2302	13.28	20,221	14.20	20,221	13.99
Men	1446	7.37	1020	9.87	1918	10.72	1162	10.11	766	10.17	6312	9.44	6312	9.53
Women	1840	9.71	3115	22.85	4835	24.13	2583	19.73	1536	15.67	13,909	18.42	13,909	18.32
Diabetes														
Both sexes	221	0.57	724	3.02	4181	11.02	6897	28.05	7872	45.41	19,895	13.97	19,895	13.78
Men	108	0.55	241	2.33	2091	11.69	3658	31.82	3723	49.42	9821	14.69	9821	14.94
Women	113	0.60	483	3.54	2090	10.43	3239	24.74	4149	42.32	10,074	13.34	10,074	12.82
Arrhythmias														
Both sexes	750	1.95	1689	7.05	3874	10.21	4403	17.91	7988	46.08	18,704	13.14	18,704	13.03
Men	348	1.78	584	5.65	1574	8.80	2227	19.37	3826	50.79	8559	12.80	8559	13.21
Women	402	2.12	1105	8.11	2300	11.48	2176	16.62	4162	42.46	10,145	13.43	10,145	13.05
Esophageal disorders														
Both sexes	1462	3.79	1260	5.26	3973	10.48	3973	16.16	4117	23.75	14,785	10.38	14,785	10.36
Men	792	4.04	499	4.83	1860	10.40	1765	15.35	1700	22.57	6616	9.89	6616	10.08
Women	670	3.54	761	5.58	2113	10.54	2208	16.87	2417	24.66	8169	10.82	8169	10.59
Asthma														
Both sexes	4141	10.74	2108	8.80	3125	8.24	1951	7.94	1424	8.21	12,749	8.95	12,749	8.88
Men	2382	12.15	716	6.93	1043	5.83	630	5.48	493	6.55	5264	7.87	5264	7.75

Continued on next page

TABLE 1. Continued

Chronic disease group	Age (y)										All ages			
	0-18		19-29		30-49		50-64		≥65		Crude ^b		Standardized ^c	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Asthma, continued														
Women	1759	9.28	1392	10.21	2082	10.39	1321	10.09	931	9.50	7485	9.91	7485	9.91
Thyroid disorders														
Both sexes	305	0.79	963	4.02	3546	9.35	3732	15.18	4283	24.71	12,829	9.01	12,829	8.87
Men	106	0.54	150	1.45	638	3.57	789	6.86	1091	14.48	2774	4.15	2774	4.27
Women	199	1.05	813	5.96	2908	14.51	2943	22.48	3192	32.56	10,055	13.32	10,055	13.00
Deficiency and other anemia														
Both sexes	868	2.25	1040	4.34	2803	7.39	2751	11.19	5148	29.70	12,610	8.86	12,610	8.75
Men	406	2.07	163	1.58	582	3.25	997	8.67	2161	28.69	4309	6.44	4309	6.65
Women	462	2.44	877	6.43	2221	11.08	1754	13.40	2987	30.47	8301	10.99	8301	10.79
Bowel disorders														
Both sexes	481	1.25	630	2.63	1843	4.86	4525	18.40	5195	29.97	12,674	8.90	12,674	8.68
Men	253	1.29	243	2.35	937	5.24	2338	20.34	2459	32.64	6230	9.32	6230	9.39
Women	228	1.20	387	2.84	906	4.52	2187	16.71	2736	27.91	6444	8.53	6444	8.09
Cancer ^d														
Both sexes	94	0.24	397	1.66	1887	4.98	3272	13.31	6334	36.54	11,984	8.42	11,984	8.28
Men	52	0.27	85	0.82	630	3.52	1483	12.90	3202	42.51	5452	8.15	5452	7.62
Women	42	0.22	312	2.29	1257	6.27	1789	13.67	3132	31.95	6532	8.65	6532	8.92
Biliary and liver disorders														
Both sexes	1141	2.96	795	3.32	3207	8.46	3392	13.80	3243	18.71	11,778	8.27	11,778	8.23
Men	599	3.05	242	2.34	1456	8.14	1459	12.69	1443	19.16	5199	7.78	5199	7.93
Women	542	2.86	553	4.06	1751	8.74	1933	14.77	1800	18.36	6579	8.71	6579	8.53
Obstructive pulmonary disorders														
Both sexes	1738	4.51	1132	4.72	2816	7.43	2506	10.19	3263	18.82	11,455	8.05	11,455	8.00
Men	910	4.64	352	3.41	1089	6.09	1097	9.54	1466	19.46	4914	7.35	4914	7.47
Women	828	4.37	780	5.72	1727	8.62	1409	10.76	1797	18.33	6541	8.66	6541	8.56
Ischemic heart disease														
Both sexes	164	0.43	156	0.65	1107	2.92	3084	12.54	6833	39.42	11,344	7.97	11,344	7.87
Men	88	0.45	102	0.99	631	3.53	1895	16.48	3513	46.64	6229	9.32	6229	9.60
Women	76	0.40	54	0.40	476	2.38	1189	9.08	3320	33.87	5115	6.77	5115	6.45

^aNumbers to the left of the prevalence figure indicate the actual number of cases observed. Prevalence can be computed by dividing the number of cases by the following corresponding denominators (and multiplying by 100); Denominators for men and women combined: 0-18 = 38,558; 19-29 = 23,968; 30-49 = 37,927; 50-64 = 24,588; ≥65 = 17,336. Denominators for men: 0-18 = 19,611; 19-29 = 10,337; 30-49 = 17,888; 50-64 = 11,496; ≥65 = 7533. Denominators for women: 0-18 = 18,947; 19-29 = 13,631; 30-49 = 20,039; 50-64 = 13,092; ≥65 = 9803.

^bA crude prevalence was computed by dividing cases observed across all ages by the total population.

^cOverall prevalence for men and women combined was standardized by age and sex; overall prevalence for men and women separately was standardized only by age (direct standardization using the 2000 US Census population).

^dPrevalence for men excluded female cancers (eg, ovarian, uterine), and prevalence for women excluded male cancers (eg, prostate, testicular).

single conditions observed in Olmsted County were generally in agreement with US statistics (Table 2). For example, national prevalence estimates indicate that approximately 30% of the adult population is affected by hypertension, increasing from 29.9% in persons 18 years or older to 70.3% in persons 65 years or older.¹⁹ These numbers were similar to the estimated prevalence of hypertension among the adult Olmsted County population (24.7% in those 18 years or older and 70.7% in those 65 years or older). However, there were greater

differences for some other diseases. For example, the prevalence of osteoarthritis in people 65 years or older was 44.4% in Olmsted County compared to 33.6% in the total US population of the same age.²⁰

DISCUSSION

Principal Findings

Using the REP medical records linkage system, we described the prevalence of the most common medical conditions in a defined US




FIGURE 1. Prevalence (per 100 population) of the 10 most prevalent disease groups in 5 broad age categories and for all ages combined (lower right panel). Prevalence figures were age and sex standardized (when applicable). Prevalence in both sexes is shown with brown bars, prevalence in men is shown with blue bars, and prevalence in women is shown with orange bars.

population across all ages, for men and women separately, and across ethnic groups. Surprisingly, the most prevalent nonacute conditions in our community were not chronic conditions related to aging such as diabetes and heart disease but rather conditions that affect both sexes and all age groups: skin disorders, osteoarthritis and joint disorders, back problems, disorders of lipid metabolism, and upper respiratory tract disease (excluding asthma). The broad disease groups that we examined in this study are useful for describing important drivers of health care utilization that might otherwise be overlooked.

Unexpectedly, almost half of the Olmsted County population of all ages received a diagnosis of skin disorders within approximately 5 years. The skin disorders category was broad and included 19 different ICD-9 groupings (including actinic keratosis, acne, and sebaceous cysts). No single skin disorder was highly prevalent, but skin disorders in combination affected a considerable proportion of all age groups in our population. Skin disorders are not typically major drivers of disability or death but may be important determinants of health care utilization and cost. For example, many of the actinic skin issues require continued observation and therapy.²¹ New models of dermatological care delivery, such as teledermatology, should be critically explored within US health care systems to increase care efficiency and reduce health care expenditures.²² Our data suggest that such efficiencies could affect a substantial proportion of the population.

The osteoarthritis and joint disorders group was also common in our population. The ICD-9 code 719 "Other and unspecified joint disorders" was assigned 216,153 times in the study time frame. The ICD-9 code 719.4 (joint pain) accounted for most of these diagnoses (176,546; 82%). Our data suggest that resources to diagnose, treat, and prevent joint pain may be required; however, joint pain occurs for multiple reasons. Overuse and activity injuries can cause short-term pain, whereas chronic conditions such as osteoarthritis and obesity may cause long-term pain.^{23,24} The underlying etiology of the joint pain cannot be determined from the ICD-9 codes, and it will be necessary to acquire additional information to determine the exact health care burden and needs for these

patients. Our data point to the need for further study of this common problem and its causes to identify areas for intervention.

Back problems were the third most prevalent disease group. Back problems and back pain are highly prevalent in the US²⁵ and have been previously classified as the eighth most costly chronic condition in patients aged 18 to 64 years.² Management of back problems can be challenging, and Carey et al²⁶ noted that patients experience similar outcomes despite a wide variation in health care professional, type of treatment, and cost of treatment. The implementation of protocols to stratify the management of patients with back pain in the primary care setting has been shown to improve health and decrease costs.²⁷ The availability of detailed information from a complete population will allow us to study current treatments for back problems and to evaluate how these treatments compare with evidence-based guidelines.²⁸ Additionally, as with skin conditions, improved management of patients with back problems could affect a substantial proportion of the population.

Disorders of lipid metabolism was the fourth most prevalent disease group. Consistent with our observation in Olmsted County, hyperlipidemia is highly prevalent in many populations throughout the US.²⁹ Hyperlipidemia contributes to multiple chronic conditions but also offers a potential target for intervention. Current guidelines for the treatment of hyperlipidemia clearly identify groups of patients most likely to benefit from treatment.³⁰ Among patients with diabetes, telephonic management of hyperlipidemia by nurses may hold promise for improving lipid control and reducing costs.³¹

Finally, other upper respiratory tract disease (excluding asthma) was the fifth most common category in our population. Similar to skin problems, the conditions included in other upper respiratory tract disease are not considered major causes of morbidity or mortality. However, these conditions are extremely common and affect all age groups. Allergic rhinitis accounted for over half of the diagnoses in this category. Allergic rhinitis alone has been estimated to affect up to 40 million Americans, and symptoms are present for more than 4 months of each year in more than half of the affected patients.³² Additionally, direct and




FIGURE 2. Age-specific prevalence (per 100 population) of the 15 most prevalent disease groups in men (blue line) compared with women (orange line). The 15 panels are presented in decreasing order of overall age- and sex-adjusted prevalence (see Table 1).




FIGURE 3. Prevalence (per 100 population) of the 10 most prevalent disease groups by ethnic group. Prevalence figures were standardized by age and sex (when applicable). Prevalence in whites is shown with brown bars, prevalence in blacks is shown with blue bars, and prevalence in Asians is shown with orange bars.

indirect health care expenditures related to allergic rhinitis were approximately \$11.2 billion in 2005.³³ Patients with allergic rhinitis often have multiple comorbid conditions including eczema, asthma, chronic sinusitis, and nasal polyps.^{32,34} Effective treatment of the conditions included in other upper respiratory tract disease may represent an ideal opportunity to improve the health care management of a considerable proportion of the community.

Strengths and Limitations

The strengths of our study include access to data on all conditions for an entire population, across age, sex, and ethnic groups. Such data are often difficult to obtain in the US because we lack a centralized health care surveillance system. For example, Medicare data contain similar diagnosis information, but the data are largely limited to elderly persons (aged 65 years and older). Data from health insurers contain similar diagnostic information, but the populations are limited to subjects who are insured,

and insured patients may be healthier than the general population.⁷

The main limitation of our study is the inability to verify the validity of ICD-9 codes. We know from previous REP studies that codes may be assigned in error, and manual review of the medical records is often needed to ascertain whether an individual truly has the disease or condition of interest.³⁵⁻³⁹ Additionally, we may have missed patients who should have been assigned a code of interest but were not. However, because many of the diagnoses represent chronic conditions, it is likely that affected patients would be seen at least once within the 5-year period. Despite these limitations, and despite differences in the methodology used for calculating prevalence, our prevalence estimates for 10 common chronic conditions were similar to published US population estimates (Table 2). These data suggest that using ICD-9 codes stored electronically for administrative purposes may be useful to estimate prevalence rates for broad groups of diseases and to monitor the health of a given population over time at relatively low cost.

Many ICD-9 codes are nonspecific, and it is unclear whether some of these code notations (eg, for joint pain) are the first indication of an underlying pathology that might be diagnosed with additional follow-up. Therefore, these data are useful to understand why people were visiting their doctors but may not be useful to understand the etiology of specific underlying diseases.

Olmsted County, Minnesota, is home to Mayo Clinic, a tertiary referral center with an international reputation. It is possible that patients might move to the area for treatment and remain as residents of the community. It is also possible that the access to a large number of medical specialists and subspecialists could result in an increased likelihood of diagnosis of specific conditions. Finally, a larger proportion of the Olmsted County population (22%) is employed by a health care institution compared with the rest of the US (9%).^{46,47} If health care employees are more likely to visit a health care professional than those who are not employed by a health care institution, our prevalence data could be substantially higher than in the rest of the US. However, we compared the prevalence of 10 common

TABLE 2. Comparison of the Prevalence (per 100 Population) of Selected Diseases and Conditions in the Total US Population vs the Olmsted County Population^a

Disease or condition	Age stratum	US population		Olmsted County population prevalence (%) ^b
		Reference	Prevalence (%)	
Hypertension		Keenan et al, 2011 ¹⁹		
	≥18 y		29.9	24.7
	≥65 y		70.3	70.7
Mood disorders		Kessler et al, 2005 ⁴⁰		
	18-29 y		21.4	23.2
	≥60 y		11.9	23.9
Diabetes		CDC, 2011 ⁴¹		
	≥20 y		11.3	9.0 ^c
	≥65 y		26.9	23.9 ^c
Osteoarthritis		Lawrence et al, 2008 ²⁰		
	≥25 y		13.9	18.5
	≥65 y		33.6	44.4
Asthma		Akinbami et al, 2011 ⁴²		
	0-17 y		9.6	10.6
	≥18 y		7.7	8.4
Osteoporosis	≥65 y	Cheng et al, 2009 ⁴³	29.7	21.4
Prostate cancer	All ages	ACS, 2011 ⁴⁴	1.6 ^d	2.3
Breast cancer	All ages	ACS, 2011 ⁴⁴	1.7 ^d	2.2
Colon cancer	All ages	ACS, 2011 ⁴⁴	0.4 ^d	0.4
HIV infection	All ages	Minnesota DOH, 2012 ⁴⁵	0.1	0.1

^aACS = American Cancer Society; CDC = Centers for Disease Control and Prevention; DOH = Department of Health; HIV = human immunodeficiency virus; ICD-9 = International Classification of Diseases, Ninth Revision.

^bAge and sex standardized using the 2000 US Census population (direct standardization).

^cPrevalence estimates include only ICD-9 codes for diabetes, not for abnormal glucose tests.

^dPrevalence calculated using the estimated US population on July 1, 2008 (men: 149,924,604; women: 154,135,120; both sexes: 304,059,724).

conditions in Olmsted County to national prevalence figures and found similar frequencies (Table 2). These comparisons suggest that the in-migration for health care, the higher probability of diagnosis associated with a tertiary care center, and the higher frequency of health care employees in Olmsted County did not artificially inflate the prevalence of the conditions that were studied.

CONCLUSION

In this article, we report the prevalence of 47 broad categories of nonacute conditions across all age groups, in men and women separately, and across ethnic groups in the Olmsted County population. The data provide insight

into current health care use in a defined US population and may predict future health care service and work force needs as well as opportunities for prevention. Finding that skin and back problems are major drivers of health care utilization affirms the importance of moving beyond the commonly recognized health care priorities such as diabetes, heart disease, or cancer. Our findings highlight opportunities to improve health care and decrease costs related to common nonacute conditions as we move forward through the changing health care landscape.

ACKNOWLEDGEMENTS

We thank Lori Klein for assistance with manuscript preparation.

SUPPLEMENTAL ONLINE MATERIAL

Supplemental material can be found online at <http://www.mayoclinicproceedings.org>.

Abbreviations and Acronyms: CCC = clinical classification code; ICD-9 = *International Classification of Diseases, Ninth Revision*; REP = Rochester Epidemiology Project

Grant Support: This study was made possible by the Rochester Epidemiology Project (grant number R01-AG034676; Principal Investigators: Walter A. Rocca, MD, MPH, and Barbara P. Yawn, MD, MSc). This project was also supported by funding from the Mayo Clinic Center for Translational Science Activities (UL1 RR024150).

Correspondence: Address to Walter A. Rocca, MD, MPH, Division of Epidemiology, Mayo Clinic, 200 First St SW, Rochester, MN 55905 (rocca@mayo.edu).

REFERENCES

- Hwang W, Weller W, Ireys H, Anderson G. Out-of-pocket medical spending for care of chronic conditions. *Health Aff (Millwood)*. 2001;20(6):267-278.
- Naessens JM, Stroebel RJ, Finnie DM, et al. Effect of multiple chronic conditions among working-age adults. *Am J Manag Care*. 2011;17(2):118-122.
- Vogeli C, Shields AE, Lee TA, et al. Multiple chronic conditions: prevalence, health consequences, and implications for quality, care management, and costs. *J Gen Intern Med*. 2007;22(suppl 3):391-395.
- Katon W, Russo J, Lin EH, et al. Cost-effectiveness of a multi-condition collaborative care intervention: a randomized controlled trial. *Arch Gen Psychiatry*. 2012;69(5):506-514.
- Hussey PS, Ridgely MS, Rosenthal MB. The PROMETHEUS bundled payment experiment: slow start shows problems in implementing new payment models. *Health Aff (Millwood)*. 2011;30(11):2116-2124.
- Bates DW, Bitton A. The future of health information technology in the patient-centered medical home. *Health Aff (Millwood)*. 2010;29(4):614-621.
- St Sauver JL, Grossardt BR, Leibson CL, Yawn BP, Melton LJ III, Rocca WA. Generalizability of epidemiological findings and public health decisions: an illustration from the Rochester Epidemiology Project. *Mayo Clin Proc*. 2012;87(2):151-160.
- St Sauver JL, Grossardt BR, Yawn BP, Melton LJ III, Rocca WA. Use of a medical records linkage system to enumerate a dynamic population over time: the Rochester Epidemiology Project. *Am J Epidemiol*. 2011;173(9):1059-1068.
- Melton LJ III. History of the Rochester Epidemiology Project. *Mayo Clin Proc*. 1996;71(3):266-274.
- Agency for Healthcare Research and Quality. Medical Expenditure Panel Survey HC-120, Appendix 3: Clinical Classification Code to ICD-9-CM Code Crosswalk. MEPS website. http://meps.ahrq.gov/mepsweb/data_stats/download_data/pufs/h120/h120_icd9codes.shtml. Accessed February 7, 2011.
- Agency for Healthcare Research and Quality. Healthcare Cost and Utilization Project: Clinical Classifications Software (CCS) for ICD-9-CM. HCUP website. www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp. Updated August 30, 2012. Accessed February 1, 2012.
- Mukherjee B, Ou HT, Wang F, Erickson SR. A new comorbidity index: the health-related quality of life comorbidity index. *J Clin Epidemiol*. 2011;64(3):309-319.
- Porta MS, International Epidemiological Association. *A Dictionary of Epidemiology*. 5th ed. Oxford, England: Oxford University Press; 2008.
- Anderson DW, Mantel N. On epidemiologic surveys. *Am J Epidemiol*. 1983;118(5):613-619.
- Deming WE. Boundaries of statistical inference. In: Johnson NL, Smith H, eds. *New Developments in Survey Sampling*. New York: Wiley-Interscience; 1969:652-670.
- Rocca WA, Cha RH, Waring SC, Kokmen E. Incidence of dementia and Alzheimer's disease: a reanalysis of data from Rochester, Minnesota, 1975-1984. *Am J Epidemiol*. 1998;148(1):51-62.
- Small area income and poverty estimates: state and county estimates for 2009. United States Census Bureau website. <http://www.census.gov/did/www/saie/data/statecounty/data/2009.html>. Updated November 29, 2011. Accessed August 1, 2012.
- US Census Bureau. 2010 Demographic Profile Data, Olmsted County, MN. <http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?src=bkrmk>. Accessed November 21, 2012.
- Keenan NL, Rosendorf KA; Centers for Disease Control and Prevention (CDC). Prevalence of hypertension and controlled hypertension—United States, 2005-2008. *MMWR Surveill Summ*. 2011;60(suppl):94-97.
- Lawrence RC, Felson DT, Helmick CG, et al; National Arthritis Data Workgroup. Estimates of the prevalence of arthritis and other rheumatic conditions in the United States: Part II. *Arthritis Rheum*. 2008;58(1):26-35.
- Ko CJ. Actinic keratosis: facts and controversies. *Clin Dermatol*. 2010;28(3):249-253.
- van der Heijden JP, de Keizer NF, Bos JD, Spuls PI, Witkamp L. Teledermatology applied following patient selection by general practitioners in daily practice improves efficiency and quality of care at lower cost. *Br J Dermatol*. 2011;165(5):1058-1065.
- Zhang Y, Jordan JM. Epidemiology of osteoarthritis. *Clin Geriatr Med*. 2010;26(3):355-369.
- Sowers MR, Karvonen-Gutierrez CA. The evolving role of obesity in knee osteoarthritis. *Curr Opin Rheumatol*. 2010;22(5):533-537.
- Deyo RA, Mirza SK, Martin BI. Back pain prevalence and visit rates: estimates from U.S. national surveys, 2002. *Spine (Phila PA 1976)*. 2006;31(23):2724-2727.
- Carey TS, Garrett J, Jackman A, McLaughlin C, Fryer J, Smucker DR. The outcomes and costs of care for acute low back pain among patients seen by primary care practitioners, chiropractors, and orthopedic surgeons: the North Carolina Back Pain Project. *N Engl J Med*. 1995;333(14):913-917.
- Hill JC, Whitehurst DG, Lewis M, et al. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomised controlled trial. *Lancet*. 2011;378(9802):1560-1571.
- Chou R, Qaseem A, Snow V, et al; Clinical Efficacy Assessment Subcommittee of the American College of Physicians; American College of Physicians; American Pain Society Low Back Pain Guidelines Panel. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society [published correction appears in *Ann Intern Med*. 2008;148(3):247-248]. *Ann Intern Med*. 2007;147(7):478-491.
- Centers for Disease Control and Prevention (CDC). Vital signs: prevalence, treatment, and control of high levels of low-density lipoprotein cholesterol—United States, 1999-2002 and 2005-2008. *MMWR Morb Mortal Wkly Rep*. 2011;60(4):109-114.
- Grundey SM, Cleeman JI, Merz CN, et al; National Heart, Lung, and Blood Institute; American College of Cardiology Foundation; American Heart Association. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation*. 2004;110(2):227-239.
- Fischer HH, Eisert SL, Everhart RM, et al. Nurse-run, telephone-based outreach to improve lipids in people with diabetes. *Am J Manag Care*. 2012;18(2):77-84.

32. Nathan RA. The burden of allergic rhinitis. *Allergy Asthma Proc.* 2007;28(1):3-9.
33. Blaiss MS. Allergic rhinitis: direct and indirect costs. *Allergy Asthma Proc.* 2010;31(5):375-380.
34. Ker J, Hartert TV. The atopic march: what's the evidence? *Ann Allergy Asthma Immunol.* 2009;103(4):282-289.
35. Leibson CL, Brown AWW, Ransom JE, et al. Incidence of traumatic brain injury across the full disease spectrum: a population-based medical record review study. *Epidemiology.* 2011;22(6):836-844.
36. Leibson CL, Naessens JM, Brown RD, Whisnant JP. Accuracy of hospital discharge abstracts for identifying stroke. *Stroke.* 1994;25(12):2348-2355.
37. Leibson CL, Needleman J, Buerhaus P, et al. Identifying in-hospital venous thromboembolism (VTE): a comparison of claims-based approaches with the Rochester Epidemiology Project VTE cohort. *Med Care.* 2008;46(2):127-132.
38. Roger VL, Killian J, Henkel M, et al. Coronary disease surveillance in Olmsted County: objectives and methodology. *J Clin Epidemiol.* 2002;55(6):593-601.
39. Yawn BP, Wollan P, St Sauver J. Comparing shingles incidence and complication rates from medical record review and administrative database estimates: how close are they? *Am J Epidemiol.* 2011;174(9):1054-1061.
40. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication [published correction appears in *Arch Gen Psychiatry.* 2005;62(7):768]. *Arch Gen Psychiatry.* 2005;62(6):593-602.
41. 2011 National Diabetes Fact Sheet. Centers for Disease Control and Prevention website. <http://www.cdc.gov/diabetes/pubs/estimates11.htm#1>. Updated May 23, 2011. Accessed February 8, 2012.
42. Akinbami LJ, Mooman JE, Liu X. Asthma prevalence, health care use, and mortality: United States, 2005-2009. *Natl Health Stat Report.* 2011;(32):1-14.
43. Cheng H, Gary LC, Curtis JR, et al. Estimated prevalence and patterns of presumed osteoporosis among older Americans based on Medicare data. *Osteoporos Int.* 2009;20(9):1507-1515.
44. Cancer prevalence: how many people have cancer? American Cancer Society website. <http://www.cancer.org/cancer/cancerbasics/cancer-prevalence>. Updated October 18, 2011. Accessed February 2, 2012.
45. HIV/AIDS prevalence and mortality tables - 2010. Minnesota Department of Health website. <http://www.health.state.mn.us/divs/idepc/diseases/hiv/stats/pmtables2010.html#table1>. Updated April 30, 2012. Accessed February 8, 2012.
46. Occupational Employment Statistics: May 2011 metropolitan and nonmetropolitan area occupational employment and wage estimates; Rochester, MN. US Bureau of Labor Statistics website. http://www.bls.gov/oes/current/oes_40340.htm#29-0000. Updated March 27, 2012. Accessed August 1, 2012.
47. Occupational Employment Statistics: May 2011 national occupational employment and wage estimates by ownership; cross-industry, private ownership only. US Bureau of Labor Statistics website. <http://www.bls.gov/oes/current/000001.htm>. Updated March 29, 2012. Accessed August 1, 2012.